

NOVEL HUMAN SECRETED PROTEINS AND  
POLYNUCLEOTIDES ENCODING THE SAME

The present application claims the benefit of U.S.  
Provisional Application Number 60/249,044, which was filed on  
5 November 15, 2000 and is herein incorporated by reference in its  
entirety.

1. INTRODUCTION

10 The present invention relates to the discovery,  
identification, and characterization of novel human  
polynucleotides encoding proteins sharing sequence similarity  
with mammalian secreted proteins. The invention encompasses the  
described polynucleotides, host cell expression systems, the  
encoded proteins, fusion proteins, polypeptides and peptides,  
antibodies to the encoded proteins and peptides, and genetically  
15 engineered animals that either lack or over express the  
disclosed genes, antagonists and agonists of the proteins, and  
other compounds that modulate the expression or activity of the  
proteins encoded by the disclosed genes that can be used for  
diagnosis, drug screening, clinical trial monitoring, the  
20 treatment of diseases and disorders, and cosmetic or  
nutriceutical applications.

2. BACKGROUND OF THE INVENTION

Human secreted proteins and growth factors have been  
implicated in a number of biological processes and medical  
25 conditions and anomalies.

3. SUMMARY OF THE INVENTION

The present invention relates to the discovery,  
identification, and characterization of nucleotides that encode  
novel human proteins, and the corresponding amino acid sequences  
30 of these proteins. The novel human proteins (NHPs) described

for the first time herein share structural similarity with animal Wnt family proteins (SEQ ID NOS:1-5) and other animal proteins including, but not limited to, disintegrins, metalloproteinases, and other human secreted proteins. SEQ ID NOS:6-8 describe a NHP that is similar to the human protein hormones chorionic gonadotrophin and follicle stimulating hormone. The novel human sequences described herein encode alternative proteins/open reading frames (ORFs) of 433, 363, and 84 amino acids in length (see SEQ ID NOS:2, 4, and 7).

10 The invention also encompasses agonists and antagonists of the described NHPs, including small molecules, large molecules, mutant NHPs, or portions thereof, that compete with native NHP, peptides, and antibodies, as well as nucleotide sequences that can be used to inhibit the expression of the described NHPs  
15 (e.g., antisense and ribozyme molecules, and open reading frame or regulatory sequence replacement constructs) or to enhance the expression of the described NHPs (e.g., expression constructs that place the described polynucleotide under the control of a strong promoter system), and transgenic animals that express a  
20 NHP sequence, or "knock-outs" (which can be conditional) that do not express a functional NHP. Knock-out mice can be produced in several ways, one of which involves the use of mouse embryonic stem cells ("ES cells") lines that contain gene trap mutations in a murine homolog of at least one of the described NHPs. When  
25 the unique NHP sequences described in SEQ ID NOS:1-8 are "knocked-out" they provide a method of identifying phenotypic expression of the particular gene as well as a method of assigning function to previously unknown genes. In addition, animals in which the unique NHP sequences described in SEQ ID  
30 NOS:1-8 are "knocked-out" provide a unique source in which to elicit antibodies to homologous and orthologous proteins that would have been previously viewed by the immune system as "self" and therefore would have failed to elicit significant antibody

responses. To these ends, gene trapped knockout ES cells have been generated in murine homologs of the described NHPs.

Additionally, the unique NHP sequences described in SEQ ID NOS:1-8 are useful for the identification of protein coding sequence and mapping a unique gene to a particular chromosome. These sequences identify biologically verified exon splice junctions as opposed to splice junctions that may have been bioinformatically predicted from genomic sequence alone. The sequences of the present invention are also useful as additional DNA markers for restriction fragment length polymorphism (RFLP) analysis, and in forensic biology.

Further, the present invention also relates to processes for identifying compounds that modulate, i.e., act as agonists or antagonists, of NHP expression and/or NHP activity that utilize purified preparations of the described NHPs and/or NHP product, or cells expressing the same. Such compounds can be used as therapeutic agents for the treatment of any of a wide variety of symptoms associated with biological disorders or imbalances.

20

#### 4. DESCRIPTION OF THE SEQUENCE LISTING AND FIGURES

The Sequence Listing provides the sequences of the NHP ORFs encoding the described NHP amino acid sequences. SEQ ID NOS:5 and 8 describe NHP ORFs and flanking regions.

25

#### 5. DETAILED DESCRIPTION OF THE INVENTION

The NHPs described for the first time herein are novel proteins that are expressed in, *inter alia*, human brain, pituitary, cerebellum, thymus, spleen, lymph node, kidney, fetal liver, prostate, testis, thyroid, adrenal gland, salivary gland, stomach, small intestine, colon, skeletal muscle, heart, uterus, placenta, mammary gland, adipose, esophagus, bladder, cervix, rectum, pericardium, hypothalamus, ovary, fetal kidney and fetal

lung (SEQ ID NOS:1-5), and/or human fetal brain, spinal cord, thymus, lymph node, lung, kidney, testis, adrenal gland, bone marrow, stomach, small intestine, colon, uterus, placenta, mammary gland, bladder, hypothalamus, fetal kidney, fetal lung, 5 gall bladder, aorta, osteosarcoma, embryo (6, 9 and 12 weeks), embryonic carcinoma, and microvascular endothelium (SEQ ID NOS:6-8).

The present invention encompasses the nucleotides presented in the Sequence Listing, host cells expressing such nucleotides, 10 the expression products of such nucleotides, and: (a) nucleotides that encode mammalian homologs of the described gene, including the specifically described NHP, and related NHP products; (b) nucleotides that encode one or more portions of a NHP corresponding to a NHP functional domain(s), and the 15 polypeptide products specified by such nucleotide sequences, including, but not limited to, the novel regions of any active domain(s); (c) isolated nucleotides that encode mutant versions, engineered or naturally occurring, of the described NHP in which all or a part of at least one domain is deleted or altered, and 20 the polypeptide products specified by such nucleotide sequences, including, but not limited to, soluble proteins and peptides in which all or a portion of the signal sequence is deleted; (d) nucleotides that encode chimeric fusion proteins containing all or a portion of a coding region of a NHP, or one of its domains 25 (e.g., a receptor or ligand binding domain, accessory protein/self-association domain, etc.) fused to another peptide or polypeptide; or (e) therapeutic or diagnostic derivatives of the described polynucleotides such as oligonucleotides, antisense polynucleotides, ribozymes, dsRNA, or gene therapy 30 constructs comprising a sequence first disclosed in the Sequence Listing. As discussed above, the present invention includes the human DNA sequences presented in the Sequence Listing (and vectors comprising the same) and additionally contemplates any

nucleotide sequence encoding a contiguous NHP open reading frame (ORF) that hybridizes to a complement of a DNA sequence presented in the Sequence Listing under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO<sub>4</sub>, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1x SSC/0.1% SDS at 68°C (Ausubel et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & Sons, Inc., New York, at p. 2.10.3) and encodes a functionally equivalent expression product. Additionally contemplated are any nucleotide sequences that hybridize to the complement of the DNA sequence that encode and express an amino acid sequence presented in the Sequence Listing under moderately stringent conditions, e.g., washing in 0.2x SSC/0.1% SDS at 42°C (Ausubel et al., 1989, *supra*), yet still encode a functionally equivalent NHP product. Functional equivalents of a NHP include naturally occurring NHPs present in other species and mutant NHPs whether naturally occurring or engineered (by site directed mutagenesis, gene shuffling, directed evolution as described in, for example, U.S. Patent No. 5,837,458 herein incorporated by reference). The invention also includes degenerate nucleic acid variants of the disclosed NHP polynucleotide sequence.

Additionally contemplated are polynucleotides encoding NHP ORFs, or their functional equivalents, encoded by polynucleotide sequences that are about 99, 95, 90, or about 85 percent similar or identical to corresponding regions of the nucleotide sequences of the Sequence Listing (as measured by BLAST sequence comparison analysis using, for example, the GCG sequence analysis package using standard default settings).

The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the described NHP gene nucleotide sequences. Such hybridization conditions may be highly stringent or less

highly stringent, as described above. In instances where the nucleic acid molecules are deoxyoligonucleotides ("DNA oligos"), such molecules are generally about 16 to about 100 bases long, or about 20 to about 80, or about 34 to about 45 bases long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first disclosed in the Sequence Listing. Such oligonucleotides can be used in conjunction with the polymerase chain reaction (PCR) to screen libraries, isolate clones, and prepare cloning and sequencing templates, etc.

Alternatively, such NHP oligonucleotides can be used as hybridization probes for screening libraries, and assessing gene expression patterns (particularly using a micro array or high-throughput "chip" format). Additionally, a series of the described NHP oligonucleotide sequences, or the complements thereof, can be used to represent all or a portion of the described NHP sequences. An oligonucleotide or polynucleotide sequence first disclosed in at least a portion of one or more of the sequences of SEQ ID NOS:1-8 can be used as a hybridization probe in conjunction with a solid support matrix/substrate (resins, beads, membranes, plastics, polymers, metal or metallized substrates, crystalline or polycrystalline substrates, etc.). Of particular note are spatially addressable arrays (*i.e.*, gene chips, microtiter plates, etc.) of oligonucleotides and polynucleotides, or corresponding oligopeptides and polypeptides, wherein at least one of the biopolymers present on the spatially addressable array comprises an oligonucleotide or polynucleotide sequence first disclosed in at least one of the sequences of SEQ ID NOS:1-8, or an amino acid sequence encoded thereby. Methods for attaching biopolymers to, or synthesizing biopolymers on, solid support matrices, and conducting binding studies thereon are disclosed in, *inter alia*, U.S. Patent Nos. 5,700,637, 5,556,752,

5,744,305, 4,631,211, 5,445,934, 5,252,743, 4,713,326,  
5,424,186, and 4,689,405 the disclosures of which are herein  
incorporated by reference in their entirety.

Addressable arrays comprising sequences first disclosed in  
5 SEQ ID NOS:1-8 can be used to identify and characterize the  
temporal and tissue specific expression of a gene. These  
addressable arrays incorporate oligonucleotide sequences of  
sufficient length to confer the required specificity, yet be  
within the limitations of the production technology. The length  
10 of these probes is within a range of between about 8 to about  
2000 nucleotides. Preferably the probes consist of 60  
nucleotides and more preferably 25 nucleotides from the  
sequences first disclosed in SEQ ID NOS:1-8.

For example, a series of the described oligonucleotide  
15 sequences, or the complements thereof, can be used in chip  
format to represent all or a portion of the described sequences.  
The oligonucleotides, typically between about 16 to about 40 (or  
any whole number within the stated range) nucleotides in length  
can partially overlap each other and/or the sequence may be  
20 represented using oligonucleotides that do not overlap.  
Accordingly, the described polynucleotide sequences shall  
typically comprise at least about two or three distinct  
oligonucleotide sequences of at least about 8 nucleotides in  
length that are each first disclosed in the described Sequence  
25 Listing. Such oligonucleotide sequences can begin at any  
nucleotide present within a sequence in the Sequence Listing and  
proceed in either a sense (5'-to-3') orientation vis-a-vis the  
described sequence or in an antisense orientation.

Microarray-based analysis allows the discovery of broad  
30 patterns of genetic activity, providing new understanding of  
gene functions and generating novel and unexpected insight into  
transcriptional processes and biological mechanisms. The use of  
addressable arrays comprising sequences first disclosed in SEQ

ID NOS:1-8 provides detailed information about transcriptional changes involved in a specific pathway, potentially leading to the identification of novel components or gene functions that manifest themselves as novel phenotypes.

5 Probes consisting of sequences first disclosed in SEQ ID NOS:1-8 can also be used in the identification, selection and validation of novel molecular targets for drug discovery. The use of these unique sequences permits the direct confirmation of drug targets and recognition of drug dependent changes in gene  
10 expression that are modulated through pathways distinct from the drugs intended target. These unique sequences therefore also have utility in defining and monitoring both drug action and toxicity.

As an example of utility, the sequences first disclosed in  
15 SEQ ID NOS:1-8 can be utilized in microarrays or other assay formats, to screen collections of genetic material from patients who have a particular medical condition. These investigations can also be carried out using the sequences first disclosed in SEQ ID NOS:1-8 *in silico* and by comparing previously collected  
20 genetic databases and the disclosed sequences using computer software known to those in the art.

Thus the sequences first disclosed in SEQ ID NOS:1-8 can be used to identify mutations associated with a particular disease and also as a diagnostic or prognostic assay.

25 Although the presently described sequences have been specifically described using nucleotide sequence, it should be appreciated that each of the sequences can uniquely be described using any of a wide variety of additional structural attributes, or combinations thereof. For example, a given sequence can be  
30 described by the net composition of the nucleotides present within a given region of the sequence in conjunction with the presence of one or more specific oligonucleotide sequence(s) first disclosed in the SEQ ID NOS:1-8. Alternatively, a



restriction map specifying the relative positions of restriction endonuclease digestion sites, or various palindromic or other specific oligonucleotide sequences can be used to structurally describe a given sequence. Such restriction maps, which are typically generated by widely available computer programs (e.g., the University of Wisconsin GCG sequence analysis package, SEQUENCHER 3.0, Gene Codes Corp., Ann Arbor, MI, etc.), can optionally be used in conjunction with one or more discrete nucleotide sequence(s) present in the sequence that can be described by the relative position of the sequence relative to one or more additional sequence(s) or one or more restriction sites present in the disclosed sequence.

For oligonucleotide probes, highly stringent conditions may refer, e.g., to washing in 6xSSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may encode or act as NHP gene antisense molecules useful, for example, in NHP gene regulation (for and/or as antisense primers in amplification reactions of NHP gene nucleic acid sequences). With respect to NHP gene regulation, such techniques can be used to regulate biological functions. Further, such sequences may be used as part of ribozyme and/or triple helix sequences that are also useful for NHP gene regulation.

Inhibitory antisense or double stranded oligonucleotides can additionally comprise at least one modified base moiety that is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine,

2-methyladenine, 2-methylguanine, 3-methylcytosine,  
 5-methylcytosine, N6-adenine, 7-methylguanine,  
 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil,  
 beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil,  
 5 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-  
 oxyacetic acid (v), wybutoxosine, pseudouracil, queosine,  
 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil,  
 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid  
 methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil,  
 10 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and  
 2,6-diaminopurine.

The antisense oligonucleotide can also comprise at least  
 one modified sugar moiety selected from the group including, but  
 not limited to, arabinose, 2-fluoroarabinose, xylulose, and  
 15 hexose.

In yet another embodiment, the antisense oligonucleotide  
 will comprise at least one modified phosphate backbone selected  
 from the group consisting of a phosphorothioate, a  
 phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a  
 20 phosphordiamidate, a methylphosphonate, an alkyl  
 phosphotriester, and a formacetal or any combination or analog  
 thereof.

In yet another embodiment, the antisense oligonucleotide is  
 an  $\alpha$ -anomeric oligonucleotide. An  $\alpha$ -anomeric oligonucleotide  
 25 forms specific double-stranded hybrids with complementary RNA in  
 which, contrary to the usual  $\beta$ -units, the strands run parallel  
 to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-  
 6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue  
 et al., 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-  
 30 DNA analogue (Inoue et al., 1987, FEBS Lett. 215:327-330).  
 Alternatively, double stranded RNA can be used to disrupt the  
 expression and function of a targeted NHP.

Oligonucleotides of the invention can be synthesized by standard methods known in the art, e.g., by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples,

- 5 phosphorothioate oligonucleotides can be synthesized (Stein et al., 1988, Nucl. Acids Res. 16:3209), and methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. USA 85:7448-7451), etc.

- 10 Low stringency conditions are well-known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory  
15 Manual (and periodic updates thereof), Cold Springs Harbor Press, N.Y., and Ausubel et al., 1989, *supra*.

- Alternatively, suitably labeled NHP nucleotide probes can be used to screen a human genomic library using appropriately stringent conditions or by PCR. The identification and  
20 characterization of human genomic clones is helpful for identifying polymorphisms (including, but not limited to, nucleotide repeats, microsatellite alleles, single nucleotide polymorphisms, or coding single nucleotide polymorphisms), determining the genomic structure of a given locus/allele, and  
25 designing diagnostic tests. For example, sequences derived from regions adjacent to the intron/exon boundaries of the human gene can be used to design primers for use in amplification assays to detect mutations within the exons, introns, splice sites (e.g., splice acceptor and/or donor sites), etc., that can be used in  
30 diagnostics and pharmacogenomics.

For example, the present sequences can be used in restriction fragment length polymorphism (RFLP) analysis to

identify specific individuals. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification (as generally described in U.S.

5 Patent No. 5,272,057, incorporated herein by reference). In addition, the sequences of the present invention can be used to provide polynucleotide reagents, e.g., PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for  
10 example, providing another "identification marker" (i.e., another DNA sequence that is unique to a particular individual). Actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments.

15 Further, a NHP gene homolog can be isolated from nucleic acid from an organism of interest by performing PCR using two degenerate or "wobble" oligonucleotide primer pools designed on the basis of amino acid sequences within the NHP products disclosed herein. The template for the reaction may be total  
20 RNA, mRNA, and/or cDNA obtained by reverse transcription of mRNA prepared from human or non-human cell lines or tissue known or suspected to express an allele of a NHP gene. The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the sequence of the desired NHP gene. The  
25 PCR fragment can then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment can be labeled and used to screen a cDNA library, such as a bacteriophage cDNA library. Alternatively, the labeled fragment can be used to isolate genomic clones via the screening  
30 of a genomic library.

PCR technology can also be used to isolate full length cDNA sequences. For example, RNA can be isolated, following standard procedures, from an appropriate cellular or tissue source (i.e.,

one known, or suspected, to express a NHP gene, such as, for example, testis tissue). A reverse transcription (RT) reaction can be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" using a standard terminal transferase reaction, the hybrid may be digested with RNase H, and second strand synthesis may then be primed with a complementary primer. Thus, cDNA sequences upstream of the amplified fragment can be isolated. For a review of cloning strategies that can be used, see e.g., Sambrook et al., 1989, *supra*.

A cDNA encoding a mutant NHP sequence can be isolated, for example, by using PCR. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known or suspected to be expressed in an individual putatively carrying a mutant NHP allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal sequence. Using these two primers, the product is then amplified via PCR, optionally cloned into a suitable vector, and subjected to DNA sequence analysis through methods well-known to those of skill in the art. By comparing the DNA sequence of the mutant NHP allele to that of a corresponding normal NHP allele, the mutation(s) responsible for the loss or alteration of function of the mutant NHP gene product can be ascertained.

Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of or known to carry a mutant NHP allele (e.g., a person manifesting a NHP-associated phenotype such as, for example, paralysis or palsy, nerve damage or degeneration, an inflammatory disorder, vision disorders, etc.), or a cDNA library can be constructed using RNA from a tissue known, or suspected, to express a mutant NHP allele. A

normal NHP gene, or any suitable fragment thereof, can then be labeled and used as a probe to identify the corresponding mutant NHP allele in such libraries. Clones containing mutant NHP sequences can then be purified and subjected to sequence analysis according to methods well-known to those skilled in the art.

10 Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known, or suspected, to express a mutant NHP allele in an individual suspected of or known to carry such a mutant allele. In this manner, gene products made by the putatively mutant tissue can be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against a normal NHP product, as described below (for screening techniques, see, for example, Harlow and Lane, eds., 15 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor).

20 Additionally, screening can be accomplished by screening with labeled NHP fusion proteins, such as, for example, alkaline phosphatase-NHP or NHP-alkaline phosphatase fusion proteins. In cases where a NHP mutation results in an expression product with altered function (e.g., as a result of a missense or a frameshift mutation), polyclonal antibodies to NHP are likely to cross-react with a corresponding mutant NHP expression product. 25 Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well-known in the art.

The invention also encompasses (a) DNA vectors that contain any of the foregoing NHP coding sequences and/or their complements (*i.e.*, antisense); (b) DNA expression vectors that 30 contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences (for example, baculo virus as described

in U.S. Patent No. 5,869,336 herein incorporated by reference);  
(c) genetically engineered host cells that contain any of the  
foregoing NHP coding sequences operatively associated with a  
regulatory element that directs the expression of the coding  
5 sequences in the host cell; and (d) genetically engineered host  
cells that express an endogenous NHP sequence under the control  
of an exogenously introduced regulatory element (i.e., gene  
activation). As used herein, regulatory elements include, but  
are not limited to, inducible and non-inducible promoters,  
10 enhancers, operators and other elements known to those skilled  
in the art that drive and regulate expression. Such regulatory  
elements include, but are not limited to, the cytomegalovirus  
(hCMV) immediate early gene, regulatable, viral elements  
(particularly retroviral LTR promoters), the early or late  
15 promoters of SV40 or adenovirus, the *lac* system, the *trp* system,  
the TAC system, the TRC system, the major operator and promoter  
regions of phage lambda, the control regions of fd coat protein,  
the promoter for 3-phosphoglycerate kinase (PGK), the promoters  
of acid phosphatase, and the promoters of the yeast  $\alpha$ -mating  
20 factors.

The present invention also encompasses antibodies and anti-  
idiotypic antibodies (including Fab fragments), antagonists and  
agonists of a NHP, as well as compounds or nucleotide constructs  
that inhibit expression of a NHP sequence (transcription factor  
25 inhibitors, antisense and ribozyme molecules, or open reading  
frame sequence or regulatory sequence replacement constructs),  
or promote the expression of a NHP (e.g., expression constructs  
in which NHP coding sequences are operatively associated with  
expression control elements such as promoters,  
30 promoter/enhancers, etc.).

The NHP or NHP peptides, NHP fusion proteins, NHP  
nucleotide sequences, antibodies, antagonists and agonists can  
be useful for the detection of mutant NHPs or inappropriately

expressed NHPs for the diagnosis of disease. The NHP or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, host cell expression systems, antibodies, antagonists, agonists and genetically engineered cells and animals can be used for

5 screening for drugs (or high throughput screening of combinatorial libraries) effective in the treatment of the symptomatic or phenotypic manifestations of perturbing the normal function of NHP in the body. The use of engineered host cells and/or animals may offer an advantage in that such systems

10 allow not only for the identification of compounds that bind to the endogenous receptor for a NHP, but can also identify compounds that trigger NHP-mediated activities or pathways.

Finally, the NHP products can be used as therapeutics. For example, soluble derivatives such as a mature NHP, or NHP

15 peptides/domains corresponding to the NHP, NHP fusion protein products (especially NHP-Ig fusion proteins, i.e., fusions of a NHP, or a domain of a NHP, to an IgFc), NHP antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate or act on downstream

20 targets in a NHP-mediated pathway) can be used to directly treat diseases or disorders. For instance, the administration of an effective amount of soluble NHP, or a NHP-IgFc fusion protein or an anti-idiotypic antibody (or its Fab) that mimics the NHP could activate or effectively antagonize the endogenous NHP

25 receptor. Soluble NHP can also be modified by proteolytic cleavage to active peptide products (e.g., any novel peptide sequence initiating at any one of the amino acids presented in the Sequence Listing and ending at any downstream amino acid). Such products or peptides can be further subject to modification

30 such as the construction of NHP fusion proteins and/or can be derivatized by being combined with pharmaceutically acceptable agents such as, but not limited to, polyethylene glycol (PEG).



1003233-1001  
Nucleotide constructs encoding such NHP products can be used to genetically engineer host cells to express such products *in vivo*; these genetically engineered cells function as "bioreactors" in the body delivering a continuous supply of a NHP, a NHP peptide, or a NHP fusion protein to the body. Nucleotide constructs encoding a functional NHP, mutant NHPs, as well as antisense and ribozyme molecules can also be used in "gene therapy" approaches for the modulation of NHP expression. Thus, the invention also encompasses pharmaceutical formulations and methods for treating biological disorders.

Various aspects of the invention are described in greater detail in the subsections below.

#### 5.1 THE NHP SEQUENCES

The cDNA sequences and corresponding deduced amino acid sequences of the described NHPs are presented in the Sequence Listing. The NHP nucleotides were obtained by aligning cDNAs from brain and kidney mRNAs (SEQ ID NOS:1-5), or bone marrow and skeletal muscle mRNAs (SEQ ID NOS:6-8) (Edge Biosystems, Gaithersburg, MD, Clontech, Palo Alto, CA) and human genomic DNA sequence. Several polymorphisms were identified during the sequencing of SEQ ID NOS:1-5, including a G/A polymorphism at nucleotide position 416 of SEQ ID NO:1 (which results in an arg or gln being present at the corresponding amino acid (aa) position 139 of SEQ ID NO:2); a G/A polymorphism at nucleotide position 206 of SEQ ID NO:3 (which results in an arg or gln being present at the corresponding aa position 69 of SEQ ID NO:4); a C/T polymorphism at nucleotide position 993 of SEQ ID NO:1 (both of which result in the same amino acid being present at the corresponding aa position of SEQ ID NO:2); a C/T polymorphism at nucleotide position 783 of SEQ ID NO:3 (both of which result in the same amino acid being present at the corresponding aa position of SEQ ID NO:4); a C/T polymorphism at

nucleotide position 1283 of SEQ ID NO:1 (which results in a val or ala being present at corresponding aa position 428 of SEQ ID NO:2); and a C/T polymorphism at nucleotide position 1073 of SEQ ID NO:3 (which results in a val or ala being present at  
5 corresponding aa position 358 of SEQ ID NO:4). SEQ ID NOS:1-5 are apparently encoded on human chromosome 17 (see GENBANK accession no. AC019316).

SEQ ID NOS:6 and 8 apparently encode a the amino acid sequence of SEQ ID NO:7 as a single exon present in human  
10 genomic sequence on chromosome 1 or both of chromosomes 4 and 6 (see GENBANK accession nos. AC048370 and AC016488).

An additional application of the described novel human polynucleotide sequences is their use in the molecular mutagenesis/evolution of proteins that are at least partially  
15 encoded by the described novel sequences using, for example, polynucleotide shuffling or related methodologies. Such approaches are described in U.S. Patent Nos. 5,830,721 and 5,837,458, which are herein incorporated by reference in their entirety.

20 NHP gene products can also be expressed in transgenic animals. Animals of any species, including, but not limited to, worms, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, birds, goats, and non-human primates, e.g., baboons, monkeys, and chimpanzees may be used to generate NHP transgenic animals.

25 Any technique known in the art may be used to introduce a NHP transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Hoppe and Wagner, 1989, U.S. Patent No. 4,873,191); retrovirus mediated gene transfer  
30 into germ lines (Van der Putten et al., 1985, Proc. Natl. Acad. Sci. USA 82:6148-6152); gene targeting in embryonic stem cells (Thompson et al., 1989, Cell 56:313-321); electroporation of embryos (Lo, 1983, Mol. Cell. Biol. 3:1803-1814); and sperm-

mediated gene transfer (Lavitrano et al., 1989, Cell 57:717-723); etc. For a review of such techniques, see Gordon, 1989, Transgenic Animals, Intl. Rev. Cytol. 115:171-229, which is incorporated by reference herein in its entirety.

5 The present invention provides for transgenic animals that carry the NHP transgene in all their cells, as well as animals that carry the transgene in some, but not all their cells, i.e., mosaic animals or somatic cell transgenic animals. The transgene may be integrated as a single transgene or in  
10 concatamers, e.g., head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al., 1992, Proc. Natl. Acad. Sci. USA 89:6232-6236. The regulatory sequences required for such a  
15 cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

When it is desired that a NHP transgene be integrated into the chromosomal site of the endogenous NHP gene, gene targeting  
20 is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous NHP gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the  
25 endogenous NHP gene (i.e., "knockout" animals).

The transgene can also be selectively introduced into a particular cell type, thus inactivating the endogenous NHP gene in only that cell type, by following, for example, the teaching of Gu et al., 1994, Science, 265:103-106. The regulatory  
30 sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant NHP gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to  
5 assay whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques that include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization  
10 analysis, and RT-PCR. Samples of NHP gene-expressing tissue, may also be evaluated immunocytochemically using antibodies specific for the NHP transgene product.

## 5.2 NHPS AND NHP POLYPEPTIDES

15 The described NHPs, NHP polypeptides, NHP peptide fragments, mutated, truncated, or deleted forms of the NHPs, and/or NHP fusion proteins can be prepared for a variety of uses. These uses include, but are not limited to, the generation of antibodies, as reagents in diagnostic assays, for  
20 the identification of other cellular gene products related to a NHP, as reagents in assays for screening for compounds that can be used as pharmaceutical reagents useful in the therapeutic treatment of mental, biological, or medical disorders and disease. Given the similarity information and expression data,  
25 the described NHPs can be targeted (by drugs, oligos, antibodies, etc.) in order to treat disease, or to therapeutically augment the efficacy of therapeutic agents.

The Sequence Listing discloses the amino acid sequences encoded by the described NHP sequences. Bioinformatics analysis  
30 reveals that the NHPs are similar to, for example Wnt-family proteins (SEQ ID NOS:1-5), or human protein hormones (SEQ ID NOS:6-8). The NHPs display initiator methionines in DNA sequence contexts consistent with translation initiation sites,

and SEQ ID NO:7 displays a hydrophobic leader sequences similar to those often found in secreted proteins. SEQ ID NO:7 also displays a predicted cleavage site at or around amino acid positions 25 or 26 that indicate the approximate position of the N-terminus of the processed, or "mature," form of the protein after cleavage by eucaryotic secretion machinery.

The NHP amino acid sequences of the invention include the amino acid sequences presented in the Sequence Listing as well as analogues and derivatives thereof. Further, corresponding NHP homologues from other species are encompassed by the invention. In fact, any NHP product encoded by the NHP nucleotide sequences described above are within the scope of the invention, as are any novel polynucleotide sequences encoding all or any novel portion of an amino acid sequence presented in the Sequence Listing. The degenerate nature of the genetic code is well-known, and, accordingly, each amino acid presented in the Sequence Listing, is generically representative of the well-known nucleic acid "triplet" codon, or in many cases codons, that can encode the amino acid. As such, as contemplated herein, the amino acid sequences presented in the Sequence Listing, when taken together with the genetic code (see, for example, Table 4-1 at page 109 of "Molecular Cell Biology", 1986, J. Darnell et al. eds., Scientific American Books, New York, NY, herein incorporated by reference) are generically representative of all the various permutations and combinations of nucleic acid sequences that can encode such amino acid sequences.

The invention also encompasses proteins that are functionally equivalent to the NHP encoded by the presently described nucleotide sequences as judged by any of a number of criteria, including, but not limited to, the ability to bind and cleave a substrate of a NHP, or the ability to effect an identical or complementary downstream pathway, or a change in

cellular metabolism (e.g., proteolytic activity, ion flux, tyrosine phosphorylation, etc.). Such functionally equivalent NHP proteins include, but are not limited to, additions or substitutions of amino acid residues within the amino acid sequence encoded by the NHP nucleotide sequences described above, but that result in a silent change, thus producing a functionally equivalent expression product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

A variety of host-expression vector systems can be used to express the NHP nucleotide sequences of the invention. Where, as in the present instance, the NHP peptide or polypeptide is thought to be a soluble or secreted molecule, the peptide or polypeptide can be recovered from the culture media. Such expression systems also encompass engineered host cells that express a NHP, or functional equivalent, *in situ*. Purification or enrichment of a NHP from such expression systems can be accomplished using appropriate detergents and lipid micelles and methods well-known to those skilled in the art. However, such engineered host cells themselves may be used in situations where it is important not only to retain the structural and functional characteristics of the NHP, but to assess biological activity, e.g., in drug screening assays.

The expression systems that may be used for purposes of the invention include, but are not limited to, microorganisms such

as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing NHP nucleotide sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing NHP nucleotide sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing NHP sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing NHP nucleotide sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the NHP product being expressed. For example, when a large quantity of such a protein is to be produced for the generation of pharmaceutical compositions of or containing NHP, or for raising antibodies to a NHP, vectors that direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited to, the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which a NHP coding sequence may be ligated individually into the vector in frame with the *lacZ* coding region so that a fusion protein is produced; pIN vectors (Inouye and Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke and Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors (Pharmacia or American Type Culture Collection) can also be used to express foreign polypeptides as fusion proteins

with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The PGEX vectors are  
5 designed to include thrombin or factor Xa protease cleavage sites so that the cloned target expression product can be released from the GST moiety.

In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express  
10 foreign polynucleotide sequences. The virus grows in *Spodoptera frugiperda* cells. A NHP coding sequence can be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).  
15 Successful insertion of NHP coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera*  
20 *frugiperda* cells in which the inserted sequence is expressed (e.g., see Smith et al., 1983, J. Virol. 46:584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems can be utilized. In cases where an adenovirus is used  
25 as an expression vector, the NHP nucleotide sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric sequence may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination.  
30 Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing a NHP product in infected hosts



(e.g., see Logan and Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient translation of inserted NHP nucleotide sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire NHP gene or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of a NHP coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (See Bitter et al., 1987, Methods in Enzymol. 153:516-544).

In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the expression product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and expression products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells that possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the expression product may be used. Such mammalian host cells

include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular, human cell lines.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines that stably express the NHP sequences described above can be engineered. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci, which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines that express the NHP product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the NHP product.

A number of selection systems may be used, including, but not limited to, the herpes simplex virus thymidine kinase (Wigler et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska and Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy et al., 1980, Cell 22:817) genes, which can be employed in tk<sup>-</sup>, hgp<sup>-</sup> or apr<sup>-</sup> cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., 1980, Proc. Natl. Acad. Sci. USA 77:3567; O'Hare et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to

mycophenolic acid (Mulligan and Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., 1981, J. Mol. Biol. 150:1); and hygromycin (Santerre et al., 1984, Gene 30:147).

Alternatively, any fusion protein can be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972-8976). In this system, the sequence of interest is subcloned into a vaccinia recombination plasmid such that the sequence's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni<sup>2+</sup>-nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

Also encompassed by the present invention are fusion proteins that direct the NHP to a target organ and/or facilitate transport across the membrane into the cytosol. Conjugation of NHPs to antibody molecules or their Fab fragments could be used to target cells bearing a particular epitope. Attaching the appropriate signal sequence to the NHP would also transport the NHP to the desired location within the cell. Alternatively targeting of NHP or its nucleic acid sequence might be achieved using liposome or lipid complex based delivery systems. Such technologies are described in "Liposomes: A Practical Approach", New, R.R.C., ed., Oxford University Press, New York and in U.S. Patent Nos. 4,594,595, 5,459,127, 5,948,767 and 6,110,490 and their respective disclosures, which are herein incorporated by reference in their entirety. Additionally embodied are novel protein constructs engineered in such a way that they facilitate

transport of the NHP to the target site or desired organ, where they cross the cell membrane and/or the nucleus where the NHP can exert its functional activity. This goal may be achieved by coupling of the NHP to a cytokine or other ligand that provides targeting specificity, and/or to a protein transducing domain (see generally U.S. Provisional Patent Application Ser. Nos. 60/111,701 and 60/056,713, both of which are herein incorporated by reference, for examples of such transducing sequences) to facilitate passage across cellular membranes and can optionally be engineered to include nuclear localization.

### 5.3 ANTIBODIES TO NHP PRODUCTS

Antibodies that specifically recognize one or more epitopes of a NHP, or epitopes of conserved variants of a NHP, or peptide fragments of a NHP are also encompassed by the invention. Such antibodies include, but are not limited to, polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')<sub>2</sub> fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

The antibodies of the invention may be used, for example, in the detection of NHP in a biological sample and may, therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal amounts of NHP. Such antibodies may also be utilized in conjunction with, for example, compound screening schemes for the evaluation of the effect of test compounds on expression and/or activity of a NHP expression product. Additionally, such antibodies can be used in conjunction gene therapy to, for example, evaluate the normal and/or engineered NHP-expressing cells prior to their introduction into the patient. Such antibodies may additionally be used as a method for the inhibition of abnormal NHP activity.

Thus, such antibodies may, therefore, be utilized as part of treatment methods.

For the production of antibodies, various host animals may be immunized by injection with the NHP, an NHP peptide (e.g., one corresponding to a functional domain of an NHP), truncated NHP polypeptides (NHP in which one or more domains have been deleted), functional equivalents of the NHP or mutated variant of the NHP. Such host animals may include, but are not limited to, pigs, rabbits, mice, goats, and rats, to name but a few.

10 Various adjuvants may be used to increase the immunological response, depending on the host species, including, but not limited to, Freund's adjuvant (complete and incomplete), mineral salts such as aluminum hydroxide or aluminum phosphate, chitosan, surface active substances such as lysolecithin, 15 pluronic polyols, polyanions, peptides, oil emulsions, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Alternatively, the immune response could be enhanced by combination and or coupling with molecules such as keyhole limpet hemocyanin, tetanus 20 toxoid, diphtheria toxoid, ovalbumin, cholera toxin or fragments thereof. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of the immunized animals.

Monoclonal antibodies, which are homogeneous populations of 25 antibodies to a particular antigen, can be obtained by any technique that provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique (Kohler and Milstein, 1975, Nature 256:495-497; and U.S. Patent No. 4,376,110), the human 30 B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal Antibodies And Cancer Therapy, Alan R. Liss,

Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated *in vitro* or *in vivo*. Production of high titers of mAbs *in vivo* makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison *et al.*, 1984, Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger *et al.*, 1984, Nature, 312:604-608; Takeda *et al.*, 1985, Nature, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. Such technologies are described in U.S. Patent Nos. 6,075,181 and 5,877,397 and their respective disclosures, which are herein incorporated by reference in their entirety. Also encompassed by the present invention is the use of fully humanized monoclonal antibodies as described in U.S. Patent No. 6,150,584 and respective disclosures, which are herein incorporated by reference in their entirety.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778; Bird, 1988, Science 242:423-426; Huston *et al.*, 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward *et al.*, 1989, Nature 341:544-546) can be adapted to produce single chain antibodies against NHP expression products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments that recognize specific epitopes may be generated by known techniques. For example, such fragments

include, but are not limited to: F(ab')<sub>2</sub> fragments, which can be produced by pepsin digestion of the antibody molecule; and Fab fragments, which can be generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression  
5 libraries may be constructed (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Antibodies to a NHP can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" a given NHP, using  
10 techniques well-known to those skilled in the art (see, e.g., Greenspan and Bona, 1993, FASEB J. 7:437-444; and Nissinoff, 1991, J. Immunol. 147:2429-2438). For example antibodies that bind to a NHP domain and competitively inhibit the binding of NHP to its cognate receptor can be used to generate anti-  
15 idiotypes that "mimic" the NHP and, therefore, bind and activate or neutralize a receptor. Such anti-idiotypic antibodies or Fab fragments of such anti-idiotypes can be used in therapeutic regimens involving a NHP signaling pathway.

Additionally given the high degree of relatedness of  
20 mammalian NHPs, the presently described knock-out mice (having never seen NHP, and thus never been tolerized to NHP) have a unique utility, as they can be advantageously applied to the generation of antibodies against the disclosed mammalian NHP (i.e., NHP will be immunogenic in NHP knock-out animals).

25 The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the  
30 invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims. All cited publications, patents,

